



Name: M. Umer farooq | Quiz Subject: Chemistry

Time Remaining: 45/45 (Minutes)

Q.1

Test 3 Gases

CHEMISTRY NMDCAT

To develop non-ideality in gases the high pressure is required:

- a. to increase the intramolecular distances
- b. to decrease the attractive forces
- c. to bring the molecules close to each other
- d. to increase the Kinetic energy of molecules

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

OB OC OD

Next

Ci









Time Remaining: 44/45 (Minutes)

Q.2

Test 3 Gases

Which one of the following parameters is very important for the non-ideal behavior of gases?

- a. low pressure and low volume
- b. high volume and high temperature
- c. low temperature and high pressure
- d. high temperature, high volume and high pressure

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

orrect Answer:

Ci







Next



7 11 11 4.9 189

Chemistry

Time Remaining: 44/45 (Minutes)

Q.3

Test 3 Gases

CHEMISTRY NMDCAT

Non-polar gases of are thought to have in them as compared to ideal ones:

- a. strong polarizability
- b. weak polarizability
- c. highl temperature
- d. low temperature

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

correct Answer:

B OC OD

Next

Back



7 11 11 10.8 189

Chemistry

Time Remaining: 44/45 (Minutes)

Q.4

Test 3 Gases

CHEMISTRY NMDCAT

With the increase in polarizability, the ideal behavior of non-polar gases:

- a. increases
- b. decreases
- c. does not change
- d. may or may not be changed

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc







Next











Time Remaining: 44/45 (Minutes)

Q.5

Test 3 Gases

CHEMISTRY NMDCAT

Ideal gasses have all the following characteristics except:

- a. The molecules occupy no space
- b. Collisions among the molecules of an ideal gas are perfectly elastic
- c. Absence of intermolecular forces
- d. All of the above are correct

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

B OC OD

Next

Back

Ci











Time Remaining: 44/45 (Minutes)

Q.6

Test 3 Gases

CHEMISTRY NMDCAT

The value of compressibility factor for an ideal gas is:

- a. zero
- b. unity
- c. fraction
- d. none of given

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next



Time Remaining: 43/45 (Minutes)

Q.7

Test 3 Gases

CHEMISTRY NMDCAT

When a graph is plotted between pressure on xaxis and the PV/RT on y-axis for an ideal gas, then:

- a. Hyperbolic curve is obtained
- b. A straight line is obtained running inbetween x-axis and y-axis
- c. A straight line is obtained running parallel to pressure axis
- d. A peak is obtained running parallel to compressibility factor axis

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

B OC OD

Next

Back

Ci











Time Remaining: 43/45 (Minutes)

Q.8

Test 3 Gases

CHEMISTRY NMDCAT

Which one of the following gases show more deviation from ideal behavior?

a. CO2

b. H_2

 $c.N_2$

d. He

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc








Next







Time Remaining: 43/45 (Minutes)



Click Here if Image Doesn't Load

Correct Answer: B C D

Next

Back







Time Remaining: 43/45 (Minutes)

Q.10

Test 3 Gases

The deciding criteria for non-ideality is:

- a. Polarity
- c. bonds
- b. Molar mass
- d. none of these

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next







Time Remaining: 43/45 (Minutes)

Q.11

Test 3 Gases

Charle's law is an example of process:

- a. isothermal
- b. isobaric
- c. isochoric
- d. All of given

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next



₹ ...| ...| 3.6 (88)

Chemistry

Time Remaining: 43/45 (Minutes)

Q.12 **Test 3 Gases** CHEMISTRY NMDCAT Boyle's law is an example of process: a. isothermal b. isobaric d. All of given c. isochoric

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

B C D

Next

Back









Time Remaining: 42/45 (Minutes)

Q.13 **Test 3 Gases** CHEMISTRY NMDCAT Which gas is likely to be the most Non - ideal? b. Cl₂ a. H₂S d. Ne c. NH₃

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

B C D

Next

Back



₹ ...| ...| 6.5 (88)

Chemistry

Time Remaining: 42/45 (Minutes)

Q.14 **Test 3 Gases** CHEMISTRY NMDCAT In Boyle's law _____ graph is more informative: a. Direct b. Inverse c. Both of the d. None of these STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer: B C D

Next

Back







Time Remaining: 42/45 (Minutes)

Q.15

Test 3 Gases

CHEMISTRY NMDCAT

If pressure remains constant at given temp the volume of an ideal gas is doubled as compared to volume at O°C:

a. -273°C

b. 273K

c. 273°C

d. 546°C

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

Ci









Next



₹ ...| ...| 1.8 K/s 188

Chemistry

Time Remaining: 42/45 (Minutes)

Q.16

Test 3 Gases

CHEMISTRY NMDCAT

Density of gas in S.I. units is expressed as:

- a. $Kg m^{-3}$
- b. $Kgdm^{-3}$
- c. $g dm^{-3}$
- d. $g m^{-3}$

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc

B C D

Next





Time Remaining: 42/45 (Minutes)

Q.17

Test 3 Gases

CHEMISTRY NMDCAT

A real gas obeying Vander Waal's equation will resemble ideal gas if

- a. Both 'a' and 'b' are large
- b. b. both 'a' and 'b' are small
- c. 'a' is small and 'b' is large
- d. 'a' is large and 'b' is s mall

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

orrect Answer:

B OC OD

Next

Back

Ci



7 11 11 4.8 188

Chemistry

Time Remaining: 42/45 (Minutes)

Q.18

Test 3 Gases

CHEMISTRY NMDCAT

Which is correct?

- a. 1 mm Hg = 1 torr = 1 atm
- b. 1mm Hg = 760 torr = 1 atm
- c. 760 mm Hg = 760 torr = 1 atm
- d. 760 mm Hg = 1 torr = 1 atm

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

correct Answer:

B OC OD

Next

Back

Ci









Time Remaining: 41/45 (Minutes)

Q.19

Test 3 Gases

CHEMISTRY NMDCAT

Among real gases the most ideal group is expected to be

- a. Halogens
- b. Noble gases
- c. Alkali metals
- d. Group VA

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc







Next









Time Remaining: 41/45 (Minutes)

Q.20

Test 3 Gases

CHEMISTRY NMDCAT

The order of rate of diffusion of gases NH_3 , H_2 , HCl and CO is:

- a. H_2 >CO >HCl > NH_3
- b. $NH_3 > H_2 > CO > HCl$
- c. $H_2 > NH_3 > CO > HCI$
- d. HCl >CO> $NH_3 > H_2$

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

orrect Answer:

Ci







Next











Time Remaining: 41/45 (Minutes)

Q.21

Test 3 Gases

CHEMISTRY NMDCAT

If $V_1 = 5$ litres for a gas at STP then what it will be at RTP?

a. >5

b. <5

c. = 5

d. 1

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next







Time Remaining: 41/45 (Minutes)

Q.22 **Test 3 Gases** CHEMISTRY NMDCAT Different gases with same K.E move with velocities b. different a. same

> d. None of these c. equal

> > STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

B OC OD

Next

Back





Time Remaining: 41/45 (Minutes)

Q.23

Test 3 Gases

CHEMISTRY NMDCAT

If two gases have same molecular mass, which statement is correct about them in H₂O:

- a. they have equal solubility in H2O at room temperature
- b. they have same b.p

()

- c. same rate of diffusion
- d. they have same number of atoms

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

OB OC OD

Next







Time Remaining: 40/45 (Minutes)

Q.24

Test 3 Gases

At absolute zero:

- a. Absolute volume is zero
- b. Gases liquefy

 \bigcirc

- c. All gases remain same
- d. Volume of gases reduce to actual volume

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

orrect Answer:

B OC OD

Next















Time Remaining: 40/45 (Minutes)

Q.25

Test 3 Gases

CHEMISTRY NMDCAT

Equal volume of H₂ and He are inserted in the same vessel. Pressure exerted by H2 and He in the ratio of:

a. 1:1

b. 2:1

c. 1:2

d. 4:1

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

correct Answer:

Ci









Next





Time Remaining: 40/45 (Minutes)

Q.26

Test 3 Gases

CHEMISTRY NMDCAT

With the increase in Vander Waal's forces the non-ideality in a gas:

- a. Increases
- b. Decreases
- c. Both a & b
- d. have no effect

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc

B OC OD

Next











Time Remaining: 40/45 (Minutes)

Q.27

Test 3 Gases

In which of the following molecule polarizability is greater

a. Cl₂

b. F₂

c. Br₂

 $d.l_2$

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next





Time Remaining: 40/45 (Minutes)

Q.28

Test 3 Gases

Correct statement about ideal gas:

- a. have forces of attraction
- b. exist in nature

Ci

- c. J. Thomson effect is not applicable
- d. all statements are incorrect

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

orrect Answer:

B OC OD

Next









Time Remaining: 39/45 (Minutes)

Q.29

Test 3 Gases

CHEMISTRY NMDCAT

The relationship between pressure and volume at same temperature is called

- a. isobar
- b. isotherm
- c. isotone
- d. isochore

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

correct Answer:

Ci









Next









Time Remaining: 39/45 (Minutes)

Q.30

Test 3 Gases

CHEMISTRY NMDCAT

At S.T.P which of the following gas has the lowest density:

a. Ar

b. Cl₂

c. CH₄

 $d. N_2$

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next



7 11 11 1.7 187

Chemistry

Time Remaining: 39/45 (Minutes)

Q.31

Test 3 Gases

CHEMISTRY NMDCAT

In gaseous state the approximate distance between molecules is how many times greater than their diameters.

a. 400

b. 600

c. 300

d. 900

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

Ci









Next



Time Remaining: 39/45 (Minutes)

Q.32

Test 3 Gases

Ammonia gas is liquefied more easily than nitrogen, hence

- a. Van der Waal's constant a & b of NH_{3 >} than of N₂
- b. Van der Waal's constant a & b of NH₃ than of N₂
- c. a(NH₃) < a(N₂) but $b(NH_3) > b(N_2)$
- d. a(NH_3) > a(N_2) but b(NH_3) < b(N_2)

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

orrect Answer:

Ci









Next











Time Remaining: 39/45 (Minutes)

Q.33

Test 3 Gases

CHEMISTRY NMDCAT

Which one of the following noble gases have high critical temperature?

a. Ne

b. Ar

c. Xe

d. Rn

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next







Time Remaining: 38/45 (Minutes)

Q.34

Test 3 Gases

CHEMISTRY NMDCAT

For an ideal gas the compressibility factor is equal to

a. 0.5

b. 1

c. 1.5

d. 2

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next











Time Remaining: 38/45 (Minutes)

Q.35

Test 3 Gases

Which of the following gases shows maximum non -ideal behavior at 0°C

a. He

b. H₂

c. N_2

d. CO2

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc







Next







Time Remaining: 38/45 (Minutes)

Q.36

Test 3 Gases

Most ideal gas at room temp is:

a. CO₂

b. NH₃

c. SO_2

 $d. N_2$

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

 \bigcirc









Next







star-portals.com/view/s



Time Remaining: 38/45 (Minutes)

Q.37

Test 3 Gases

Who studied the physical behaviour of real gases:

- a. Clausius
- b. Maxwell

 \bigcirc

- c. Boltzmann
- d. Van der Waal's

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:

B OC OD

Next

Back









star-portals.com/view/s



Time Remaining: 37/45 (Minutes)

Q.38

Test 3 Gases

CHEMISTRY NMDCAT

Gases deviate from ideal behavior at high pressure . Which is correct for non-ideality?

- a. at high pressure, the gas molecule move in one direction only
- b. at high pressure, the collision between the gas molecule are increased manifold
- c. at high pressure, the volume of gas becomes insignificant
- d. at high pressure, the intermolecular attractions become significant

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

correct Answer:

B OC OD

Next

Back

Ci

Chemistry

Time Remaining: 37/45 (Minutes)

Q.39

Test 3 Gases

CHEMISTRY NMDCAT

The density of a gas can be determined by formula:

a.
$$d = \frac{PM}{RT}$$

a.
$$d = \frac{PM}{RT}$$
 b. $d = \frac{RT}{PM}$

c.
$$d = \frac{PMR}{T}$$
 d. $d = \frac{PMT}{R}$

$$d = \frac{PMT}{R}$$

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:



 \bigcirc







Next

Back







star-portals.com/view/s



Time Remaining: 37/45 (Minutes)

Q.40

Test 3 Gases

The value of R (in Nm K⁻¹ mol⁻¹) is

a. 8.214

b. 8.314

c. 0.0321

d. 62.4

STAR INSTITUTE LAHORE

Click Here if Image Doesn't Load

Correct Answer:







Submit Quiz

Back



To develop non-ideality in gases the high pressure is required:

- a. to increase the intramolecular distances
- b. to decrease the attractive forces
- c. to bring the molecules close to each other
- d. to increase the Kinetic energy of molecules

THE STAR INSTITUTE

s screen'سید اختر عباس جعفری



Which one of the following parameters is very important for the non-ideal behavior of gases?

- a. low pressure and low volume
- b. high volume and high temperature
- c. low temperature and high pressure
- d. high temperature, high volume and high pressure

THE STAR INSTITUTE



Non-polar gases of are thought to have ____ in them as compared to ideal ones:

- a. strong polarizability
- b. weak polarizability
- c. highl temperature
- d. low temperature

THE STAR INSTITUTE

s screen'سید اختر عباس جعفری



With the increase in polarizability, the ideal behavior of non-polar gases:

- a. increases
- b. decreases

C

- c. does not change
- d. may or may not be changed



Ideal gasses have all the following characteristics except:

- a. The molecules occupy no space
- b. Collisions among the molecules of an ideal gas are perfectly elastic
- c. Absence of intermolecular forces
- d. All of the above are correct

C



The value of compressibility factor for an ideal gas is:

a. zero

c. fraction

C

b. unity

d. none of given



When a graph is plotted between pressure on x-axis and the PV/RT on y-axis for an ideal gas, then:

- a. Hyperbolic curve is obtained
- b. A straight line is obtained running inbetween x-axis and y-axis
- c. A straight line is obtained running parallel to pressure axis
- d. A peak is obtained running parallel to compressibility factor axis

THE STAR INSTITUTE



Which one of the following gases show more deviation from ideal behavior?

a. *CO*₂ c.*N*₂

ن

b. H_2 d. He





At what temp the gases will is likely to obey the K.M.T?

- a. low
- c. high

C

- b. highest
- d. lowest



The deciding criteria for non-ideality is:

a. Polarity

c. bonds

C

b. Molar mass

d. none of these



Charle's law is an example of process:

- a. isothermal
- c. isochoric

C

- b. isobaric
- d. All of given



Boyle's law is an example of process:

- a. isothermal
- c. isochoric

C

- b. isobaric
- d. All of given



Which gas is likely to be the most Non - ideal?

a. H₂S

c. NH₃

C

b. Cl₂

d. Ne



In Boyle's law _____ graph is more informative:

a. Direct

C

- c. Both of the
- b. Inverse
- d. None of these



If pressure remains constant at given temp the volume of an ideal gas is doubled as compared to volume at O°C:

a. -273°C

c. 273°C

C

b. 273K

d. 546°C

(273)-2



Density of gas in S.I. units is expressed as:

a. $Kg m^{-3}$ c. $g dm^{-3}$

b. $Kgdm^{-3}$ d. $g m^{-3}$

C



Density of gas in S.I. units is expressed as:

a. $Kg m^{-3}$

c. $g \ dm^{-3}$

C

b. $Kgdm^{7}$ d. $g m^{-3}$

A real gas obeying Vander Waal's equation wil resemble ideal gas if

- a. Both 'a' and 'b' are large
- b. b. both 'a' and 'b' are small
- c. 'a' is small and 'b' is large
- d. 'a' is large and 'b' is s mall

 \bigcirc



Which is correct?

 \Box

- a. 1 mm Hg = 1 torr = 1 atm
- b. 1mm Hg = 760 torr = 1 atm
- c. 760 mm Hg = 760 torr = 1 atm
- d. 760 mm Hg = 1 torr = 1 atm



Among real gases the most ideal group is expected to be

a. Halogens

C

- c. Alkali metals
- b. Noble gases
- d. Group VA



The order of rate of diffusion of gases NH_3 , H_2 , HCl and CO is:

- a. H_2 >CO >HCl > NH_3
- b. $NH_3 > H_2 > CO > HCI$
- c. $H_2 > NH_3 > CO > HCI$
- d. HCl >CO> $NH_3 > H_2$

ن



If $V_1 = 5$ litres for a gas at STP then what it will be at RTP?

a. >5

c. =5

ن

b. <5

d. 1



Different gases with same K.E move with ____ velocities

a. same

c. equal

ن

b. different

d. None of these



If two gases have same molecular mass, which statement is correct about them in H₂O:

- a. they have equal solubility in H₂O at room temperature
- b. they have same b.p

 \Box

- c. same rate of diffusion
- d. they have same number of atoms



At absolute zero:

- a. Absolute volume is zero
- b. Gases liquefy

C

- c. All gases remain same
- d. Volume of gases reduce to actual volume



Equal volume of H₂ and He are inserted in the same vessel.

Pressure exerted by H₂ and He in the ratio of:

a. 1:1 b. 2:1

 \bigcirc

c. 1:2 d. 4:1



With the increase in Vander Waal's forces the non-ideality in a gas:

- a. Increases
- c. Both a & b

ن

- b. Decreases
- d. have no effect



In which of the following molecule polarizability is greater

a. Cl₂

c. Br₂

ن

b. F₂

d. 1,



Correct statement about ideal gas:

- a. have forces of attraction
- b. exist in nature

ن

- c. J. Thomson effect is not applicable
- d. all statements are incorrect

The relationship between pressure and volume at same temperature is called

- (a) isobar
- (b) isotherm
- (c) isotone

ن

(d) isochore



At S.T.P which of the following gas has the lowest density:

(a) Ar

(b) Cl₂

(c) CH₄

(d) N₂

THE STAR INSTITUTE

s screen'سيد آختر عباس جعفری

C



In gaseous state the approximate distance between molecules is how many times greater than their diameters.

(a) 400

(b) 600

(c) 300

(d) 900

THE STAR INSTITUTE

سيد آختر عباس جعفري's screen



Ammonia gas is liquefied more easily than nitrogen, hence

- (a) Van der Waal's constant a & b of NH_{3 >} than of N
- (b) Van der Waal's constant a & b of NH_{3 <} than of N
- (c) a(NH_3) < a(N_2) but b(NH_3) > b(N_2)
- (d) a(NH_3) > a(N_2) but b(NH_3) < b(N_2)

ن



Which one of the following noble gases have high critical temperature?

(a) Ne

C

(c) Xe

(b) Ar

(d) Rn



For an ideal gas the compressibility factor is equal to

(a) 0.5

(c) 1.5

ن

(b)

(d) 2



Which of the following gases shows maximum non —ideal behavior at 0°C

(a) He

(b) H_2

(c) N_2

ن

(d) CO₂





Most ideal gas at room temp is:

(a) CO₂

(b) NH₃

(c) SO₂

ن

(d) N_2



Q. 37 Who studied the physical behaviour of real gases:

- (a) Clausius
- (b) Maxwell
- (c) Boltzmann
- (d) Van der Waal's

ن



Gases deviate from ideal behavior at high pressure . Which is correct for non-ideality?

- (a) at high pressure, the gas molecule move in one direction only
- (b) at high pressure, the collision between the gas molecule are increased manifold
- (c) at high pressure, the volume of gas becomes insignificant
- (d) at high pressure, the intermolecular attractions become significant

 \bigcirc



The density of a gas can be determined by formula:

(a)
$$d = \frac{PM}{RT}$$

C

(c)
$$d = \frac{PMR}{T}$$
 (d) $d = \frac{PMT}{R}$



The value of R (in Nm K⁻¹ mol⁻¹) is

(a)8.214

(c) 0.0321

ن

(b) 8.314

(d) 62.4